LE MONITORAGE DE STIMULATIONS OVARIENNES : DU PRÉSENT AU FUTUR

(3D AUTOMATISÉ, INTELLIGENCE ARTIFICIELLE, INTÉRÊT DU BILAN HORMONAL)

Dr Geoffroy ROBIN, MCU-PH

Service d'AMP et Préservation de la Fertilité (Dr DECANTER)

UF de Gynécologie Endocrinienne - Service de Gynécologie Médicale, Orthogénie et Sexologie (Pr CATTEAU-JONARD)

Importance relative des différents acteurs en Reproduction Humaine

La réunion de décision ou staff quotidien AMP

- Flux de patientes
- Nombre de POV, IIU, DEC/TEC réalisables/jour
- Intégrer des critères hormonaux et échographiques
- Critères de déclenchement

- Freeze-all ou « TEF » ?
- Informatisation en temps réel des tentatives
 - *Dilemme de la fiche de stim « papier »*
- Enseignement / formation
- Sérologies, RT-PCR Covid, Cs anesth & co...

Le minimum « syndical » d'un monitoring de l'ovulation

- Echographie endovaginale +++
- **La mesure du diamètre du (ou des) follicule(s) en croissance**
- **La mesure de l'épaisseur endométriale**
(au minimum au moment du déclenchement)
 - **Indiquer aspect de l'endomètre si besoin**

Les évolutions techniques...

Estimation de la croissance folliculaire

- Le diamètre moyen :
 - ✓ *moyenne des diamètres orthogonaux sur le plus grand plan du follicule.*
- Hors FIV : 1 ou 2 follicules dominants de structure sphérique
- Cas particulier de la FIV :
 - ✓ *pressions exercées par follicules en croissance adjacents → perte de la structure sphérique des follicule → struct. complexes 3D.*
 - ✓ *Analyse soigneuse et individuelle de chaque follicule pour trouver le plus grand plan de chacun d'entre-eux.*

La maturité folliculaire

A critical review of bi-dimensional and three-dimensional ultrasound techniques to monitor follicle growth: do they help improving IVF outcome?

Alberto Revelli^{1,2*}, Giorgia Martiny¹, Luisa Delle Piane², Chiara Benedetto¹, Paolo Rinaudo³ and Ilan Tur-Kaspa⁴

Author	Year	Study Design	Population	US Technique	n	US Group	% MII	% fert	% cleavage	% good quality embryo
Inaudi 1995 [25]	R	Retrospective	n.e.	n.e.	n.e.	n.e.	n.e.	Increasing from 12 mm onwards	Increasing from 12 mm onwards	None
Dubey 1995 [26]	R	Retrospective	n.e.	n.e.	n.e.	n.e.	n.e.	n.e.	n.e.	n.e.
Ectors 1997 [17]	P	Prospective	n.e.	n.e.	n.e.	n.e.	n.e.	n.e.	n.e.	n.e.
Bergh 1998 [18]	P	Prospective	33.0	Long	IVF/ICSI	n.e.	n.e.	n.e.	n.e.	n.e.
Teisser 2000 [22]	P	Prospective	28.0	Long	Only IVF	150	14	n.e.	n.e.	n.e.
Triwitayakom 2003 [14]	P	Prospective	35.9	Long short	Only ICSI	991	10-14	n.e.	n.e.	n.e.
Rosen 2008 [19]	P	Prospective	34.1	Long shortantag	IVF/ICSI	2934	10-13-16-19	n.e.	n.e.	n.e.
Lee 2010 [23]	P	Prospective	33.5	Long	Only ICSI	819	18-20	n.e.	n.e.	n.e.
Mehri 2013 [20]	P	Prospective	n.e.	Long	IVF/ICSI	360	15-18	n.e.	n.e.	n.e.

“hCG was administered when at least three follicles with a diameter > 16-18 mm were observed on ultrasound at the same time...”

Conclusions: The study of follicular size by 2D-US is of limited usefulness in helping in the identification of follicles containing the best oocytes and in choosing the best moment to trigger ovulation. Possibly the value of US in this area will be improved by large prospective studies in which automated 3D-US will be used.

From a circle to a sphere: the ultrasound imaging of ovarian follicle with 2D and 3D technology

Claudia Re^a, Mario Mignini Renzini^b, Amelia Rodriguez^c, Mariabeatrice Dal Canto^b, Matteo Buccheri^d, Sandro Sacchi^a, Silvia Bartolucci^d, Rubens Fadini^b and Antonio La Marca^{a,d}

Best setting for antral follicle count

Probe frequency: Medium
Gain: -5
SRI (Speckle reduction imaging): 5
CRI (Crossx beam): 1
CRI filter: High
Dynamic range: 8
Harmonics: Medium

Figure 2. Ovarian follides have frequently an irregular shape and more than one diameter should be measured to calculate the mean dimension.

Figure 3. The follicular measurement using sonoAVC, multiplanar view.

Monitoring automatisé 3D: quel apport?

- Acquisition rapide par balayage du « volume » ovarien
- En théorie :
 - ✓ Evaluation du diamètre moyen de tous les follicules
 - ✓ Evaluation des volumes de chacun des follicules

➔ **Plus de précision et de reproductibilité pour apprécier la maturité folliculaire?**
- En pratique :
 - ✓ le mode Sono-AVC est applicable sans retouche chez 60%-70% des patientes...
 - ✓ dans au moins 30-40% des cas : travail de l'image 3D après l'examen... parfois long et fastidieux...
- Les problèmes techniques persistants : ovaires volumineux, ovaires « lointains »...

Monitoring hors-FIV

Figure 43. Monitoring périovulatoire en induction monofolliculaire.

A. Échographie 3D en mode *Sonography-based automatic volume calculation* (SonoAVC™) : ovaires micropolykystiques sous clomifène j13 : follicule préovulatoire unique de 23 mm (en rouge) associé à de nombreux microfollicules inférieurs à 10 mm.

B. Endomètre préovulatoire en « cible » (11 mm) et glaire cervicale bien visible (flèche).

C. Doppler : corps jaune postovulatoire triangulaire avec couronne vasculaire typique.

Monitoring de la croissance folliculaire en FIV

Toute référence à cet article doit porter la mention : Ardaens Y, Robin G. Imagerie de l'hypofertilité de la femme. EMC - Radiologie et imagerie médicale - génito-urinaire - gynéco-obstétricale - mammaire 2018;13(4):1-23 [Article 34-620-E-10].

Et si on parlait de « volume folliculaire »?

Prospective evaluation of automated follicle monitoring in 58 in vitro fertilization cycles: follicular volume as a new indicator of oocyte maturity

Adela Rodríguez-Fuentes, M.D.,^a Jairo Hernández, Ph.D.,^a Rocío García-Guzmán, M.D.,^a Elena China, M.Sc.,^a Laura Iaconianni, M.D.,^b and Angela Palumbo, M.D., Ph.D.^a

TABLE 3

Relationship between the number of mature oocytes and different follicular volumes.

Follicle volume cutoff	Number of mature oocytes/number of follicles in range	Number of follicles in range/number of mature oocytes
≥ 0.8 cc	121.26%	82.47%
≥ 0.7 cc	109.07%	91.69%
≥ 0.6 cc	100.88%	98.60%
≥ 0.5 cc	94.11%	106.26%
≥ 0.4 cc	88.74%	112.69%

Note: When a cutoff of 0.6 cc is chosen, the maximum number of follicles at or above this cutoff is closest to the number of mature oocytes retrieved. Follicular volumes are those measured on the day of hCG.

Rodríguez-Fuentes. Measurement of follicle volume using SonoAVC. Fertil Steril 2010.

FIGURE 1

(a) Example of a patient with good quality of image. All follicles were picked up automatically and no postprocessing work was necessary. (b) Example of a patient with poor quality of image: note that one follicle needed manual measurement.

Rodríguez-Fuentes. Measurement of follicle volume using SonoAVC. Fertil Steril 2010.

Alors : 2D-r

Prospective cohort study of three- versus two-dimensional ultrasound for prediction of oocyte maturity

Doron Shmorgun, M.D.,^{a,b} Edward Hughes, M.D.,^{a,c} Patrick Mohide, M.D.,^{a,c} and Robin Roberts,

Human Reproduction, Vol.26, No.1 pp. 127–133, 2011
Advanced Access publication on November 26, 2010 doi:10.1093/humrep/deq320

human reproduction ORIGINAL ARTICLE Infertility

Comparison of automated and manual follicle monitoring in an unselected population of 100 women undergoing controlled ovarian stimulation

Baris Ata*, Ayse Seyhan, Shrinivasan Krishnamurti

International Journal of Gynecology and Obstetrics

Contents lists available at ScienceDirect

journal homepage: www.elsevier.com/locate/ijgo

CLINICAL ARTICLE

Three-dimensional sonography-based automated volume calculation (SonoAVC) versus two-dimensional manual follicular tracking in in vitro fertilization

Neeta Singh*, B.R. Usha, Nisha Malik, Neena Malhotra, Sangita Pant, Perumal Vanamail

Assisted Reproductive Techniques Centre, Department of Obstetrics and Gynaecology, All India Institute of Medical Sciences, New Delhi, India

Reliability of automated volumetric measurement of multiple growing follicles in controlled ovarian hyperstimulation

Frederic Lamazou, M.D., Elisangela Arbo, M.D., Samuel Salama, M.D., Ph.D., René Frydman, M.D., and Renato Fanchin, M.D.

Dr Maximilian Murtinger

monitoring of ovarian stimulation with 3D transvaginal ultrasound plus automated volume count

Dr Maximilian Murtinger received his MD at the University of Vienna in 1986. After specialization as a trauma surgeon in 2003, he changed to the field of reproductive medicine and endocrinology and is currently finishing his specialization in obstetrics and gynecology. His current interest is in ultrasound techniques, especially related to follicular measurements in stimulated cycles and in establishing clinical standards related to the quality criteria set by the European Union directive on tissues and cells (2004/23/EC). In addition, Dr Murtinger is developing requirements for software used in obstetrics and gynecology, especially for reproductive medicine and endocrinology.

➔ Automated Sono-AVC ultrasound can be used interchangeably with manual 2D USG for follicle tracking during COS giving comparable ART outcomes with the added advantage of saving time.

The value of automated follicle volume measurements in IVF/ICSI

Frank Vandekerckhove^{**†}, Victoria Bracke[†] and Petra De Sutter

Centre for Reproductive Medicine, University Hospital Ghent, Ghent, Belgium

Table 2 | Evaluation of work flow.

	Manual 2D	SonoAVC	SonoAVC with post-processing (pp.)	2D versus SonoAVC with pp.
IN COS CYCLES				
Raine-Fenning et al. (6)	236.10 ± 57.07 s	39.00 ± 6.00 s	180.50 ± 63.55 s	<i>P</i> < 0.01
Rodriguez-Fuentes et al. (17, 40)	9.6 min	–	5.6 min	4 min time gain per patient
Deb et al. (19)	324.47 ± 162.22 s	41.06 ± 11.12 s	132.05 ± 56.23 s	<i>P</i> < 0.001
Deutch et al. (31)	361 s (129–878)	–	133 s (73–271)	<i>P</i> < 0.001, 7.6 min time gain per patient
Deutch et al. (38)	56.8 s	5.8 s	–	<i>P</i> < 0.001 without pp.
Sherbahn et al. (41)	6.2 min	–	5.6 min	
IN SPONTANEOUS CYCLES				
Raine-Fenning et al. (14)	26.36 ± 4.35 s	9.34 ± 0.67 s	–	<i>P</i> < 0.01 without pp.
MEASURING tAFC				
Deb et al. (13)	71 ± 8.48 s	26 ± 2.67 s	112 ± 29.86 s	<i>P</i> < 0.001

Table 3 | Investigation of operator independence.

	Manual 2D	SonoAVC	SonoAVC with pp.	<i>P</i>
MEASURING tAFC				
Deb et al. (13)	19.26 ± 10.55* ICC = 0.979	6.51 ± 4.79*	18.42 ± 10.53* ICC = 0.997	0.006**
FOLLICLE MEASUREMENTS				
Salama et al. (29)		ICC < 0.08	ICC = 0.97	
Bouhanna et al. (43)	Intra observer variability	ICC = 0.62	ICC = 0.96	
	Inter observer variability	ICC = 0.83	ICC = 0.94	

ICC, intra class correlation coefficient; *Mean ± SD; **Comparison of manual 2D and SonoAVC with pp.

Conclusion: This overview of previously published literature shows how Sono-AVC offers advantages for clinical practice, without losing any accuracy or reliability. Doctors should be motivated to the general use of follicular volumes instead of follicular diameters.

Nr.	d(V)	dx	dy	dz	mn. d	V
	mm	mm	mm	mm	mm	cm ³
1	22.7	34.4	21.8	18.8	25.0	6.09
2	21.2	28.7	26.0	13.5	22.8	4.98
3	19.0	26.4	21.6	14.7	20.9	3.59
4	17.3	28.6	22.0	8.8	19.8	2.72
5	16.8	26.4	20.0	11.1	19.2	2.47
6	11.3	29.5	10.0	6.6	15.3	0.76
7	8.6	10.4	8.5	7.6	8.8	0.34
8	7.8	12.1	7.4	5.6	8.4	0.25
9	7.5	10.4	7.5	6.4	8.1	0.22

→ Logiciel AMP du centre → ABM
(JFIV© ou MEDIFIRST©)

Le monitoring de la croissance endométriale : la mesure de l'épaisseur endométriale

- Elle doit se faire au niveau du fond utérin (zone « en grain de café »).
- Noter l'aspect : « 3F », échogénicité...

- La mesure se fait de la jonction endomètre—myomètre antérieur jusqu'à cette même jonction postérieure **perpendiculairement** à l'axe de la cavité utérine.

Endometrial thickness and pregnancy rates after IVF: a systematic review and meta-analysis

Annemieke Kasius^{1†}, Janine G. Smit^{1†}, Helen L. Torrance¹,
 Marinus J.C. Eijkemans², Ben Willem Mol³, Brent C. Opmeer⁴,
 and Frank J.M. Broekmans¹

Table III ORs and 95% CI for clinical pregnancy at different EMT cut-off values.

Cut-off (mm)	No. of patients	OR (95% CI)	P-value	Included studies
6	6886	0.46 (0.20–1.05)	$P = 0.07$	Grant <i>et al.</i> (2007), Richter <i>et al.</i> (2007), Al-Ghamdi <i>et al.</i> (2008), Chen <i>et al.</i> (2010), Singh <i>et al.</i> (2011)
7	10 612	0.42 (0.27–0.67)	$P = 0.0003$	Basir <i>et al.</i> (2002), Richter <i>et al.</i> (2007), Al-Ghamdi <i>et al.</i> (2008), Bozdogan <i>et al.</i> (2009), Okohue <i>et al.</i> (2009), Chen <i>et al.</i> (2010), Kinay <i>et al.</i> (2010), Kuc <i>et al.</i> (2011), Zhao <i>et al.</i> (2012)
8	7247	0.56 (0.44–0.70)	$P < 0.00001$	Basir <i>et al.</i> (2002), Aboulghar <i>et al.</i> (2005), Rashidi <i>et al.</i> (2005), Richter <i>et al.</i> (2007), Al-Ghamdi <i>et al.</i> (2008), Traub <i>et al.</i> (2009), Chen <i>et al.</i> (2010), Singh <i>et al.</i> (2011)
9	8221	0.60 (0.50–0.73)	$P < 0.00001$	Dietterich <i>et al.</i> (2002), Zhang <i>et al.</i> (2005), Richter <i>et al.</i> (2007), Al-Ghamdi <i>et al.</i> (2008), Chen <i>et al.</i> (2010)
10	7446	0.68 (0.61–0.76)	$P < 0.00001$	Rinaldi <i>et al.</i> (1996), De Geyter <i>et al.</i> (2000), Dietterich <i>et al.</i> (2002), Richter <i>et al.</i> (2007), Al-Ghamdi <i>et al.</i> (2008), Chen <i>et al.</i> (2010), Singh <i>et al.</i> (2011)
11	7451	0.84 (0.61–1.15)	$P = 0.28$	De Geyter <i>et al.</i> (2000), Dietterich <i>et al.</i> (2002), Richter <i>et al.</i> (2007), Al-Ghamdi <i>et al.</i> (2008), Chen <i>et al.</i> (2010)
12	7873	0.89 (0.55–1.44)	$P = 0.65$	Dietterich <i>et al.</i> (2002), Aboulghar <i>et al.</i> (2005), Rashidi <i>et al.</i> (2005), Zhang <i>et al.</i> (2005), Richter <i>et al.</i> (2007), Al-Ghamdi <i>et al.</i> (2008), Chen <i>et al.</i> (2010); Singh <i>et al.</i> (2011)

ORs for clinical pregnancy improve with increasing thickness. However, ORs remain significantly lower up to a thickness of 10 mm.

Figure 6 Forest plot of clinical pregnancy for women with EMT ≤ 7 mm and women with EMT > 7 mm. The probability of clinical pregnancy is significantly lower for women with EMT ≤ 7 mm. The I² statistic was 41%, indicating that study heterogeneity was low.

Endometrial thickness and pregnancy rates after IVF: a systematic review and meta-analysis

Annemieke Kasius^{1†}, Janine G. Smit^{1†}, Helen L. Torrance¹,
Marinus J.C. Eijkemans², Ben Willem Mol³, Brent C. Opmeer⁴,
and Frank J.M. Broekmans¹

Figure 5 Forest plot of ongoing pregnancy and live birth for women with EMT ≤ 7 mm and EMT > 7 mm. Chances of live birth and ongoing pregnancy are lower for women with EMT ≤ 7 mm, but the results are not significant. The I² statistic of 11% indicates that study heterogeneity was low. CI, confidence interval.

- Tenir compte de l'épaisseur... mais pas que...
 - Quelques grossesses ont été rapportées sur des épaisseurs endométriales très faibles (4mm)
 - Importance de l'aspect (échogénicité)... et d'autres paramètres encore « mystérieux » (vascularisation...)?
- Quant au volume endométrial : pas plus prédictif du succès de la FIV que l'épaisseur endométriale?

ENDOMETRIAL VOLUME AND SUB-ENDOMETRIAL VASCULARITY AS PREDICTORS OF IVF SUCCESS

The measurement of endometrial volume and sub-endometrial vascularity to replace the traditional endometrial thickness as predictors of *in-vitro* fertilization success

Ahmed M. Maged, Ahmed M. Kamel, Fouad Abu-Hamila, Rasha O. Elkomy, Omar A. Ohida, Sarah M. Hassan, Radwa M. Fahmy and Wafaa Ramadan

J Ultrasound Med 2021; 9999:1-9

Table 4. Predictive value of endometrial thickness and 3d Doppler indices.

	Cutoff value	Sensitivity (%)	Spedficity (%)	PPV (%)	NPV (%)	Accuracy (%)	p-value for AUC
Endometrial thickness (mm)							
Day of triggering	10.25	52.4	54.1	28.8	76.7	49.2	.911
Day of ET	10.95	57.1	57.4	31.6	79.5	59.2	.212
Sub-endometrial VI (%)							
Day of triggering	1.56	52.4	57.1	26.2	75	56.7	.361
Day of ET	1.08	71.4	62.3	39.5	86.4	65.3	.037*
Sub-endometrial FI (0-100)							
Day of triggering	30.4	61.9	59	34.2	81.8	57.4	.315
Day of ET	29.9	61.9	54.1	31.7	80.5	56.6	.367
Sub-endometrial VFI (0-100)							
Day of triggering	0.25	81	47.5	37.4	87.9	68	.014*
Day of ET	0.25	76.2	45.9	32.7	84.8	66.7	.023*
Endometrial volume (cm ³)							
Day of triggering	3.2	81	42.6	32.7	86.7	65.3	.037*
Day of ET	3.27	66.7	52.5	32.6	82.1	64.1	.056

Diagonal segments are produced by ties.

A : thickness at hCG-D ;
 C : volume at hCG-D ;

Diagonal segments are produced by ties.

B : thickness at FET
 D : volume at FET

ESHRE guideline: ovarian stimulation for IVF/ICSI†

The ESHRE Guideline Group on Ovarian Stimulation, Ernesto Bosch¹, Simone Broer², Georg Griesinger³, Michael Grynberg⁴, Peter Humaidan⁵, Estratios Kolibianakis⁶, Michal Kunicki⁷, Antonio La Marca⁸, George Lainas⁹, Nathalie Le Clef¹⁰, Nathalie Massin¹¹, Sebastiaan Mastenbroek¹², Nikolaos Polyzos¹³, Sesh Kamal Sunkara¹⁴, Tanya Timeva¹⁵, Mira Töyli¹⁶, Janos Urbancsek¹⁷, Nathalie Vermeulen¹⁰, and Frank Broekmans^{2,*}

Recommendations

Routine monitoring of endometrial thickness during ovarian stimulation is probably not recommended.

Conditional ⊕○○○

The guideline group suggests performing a single measurement of the endometrium during ultrasound assessment on the day of triggering or oocyte pick-up to counsel patients on potential lower pregnancy chance.

GPP

L'endomètre au cours de la transition folliculo-lutéale

- De plus en plus échogène
- Attention aux utérus intermédiaires !!!

FIGURE 2

The cycles were sorted into six groups according to the extent of the upward hyperechogenic transformation of the endometrium.

Fanchin. Endometrial echogenicity and receptivity. Fertil Steril 1999.

FIGURE 3

Clinical pregnancy rates according to the extent of hyperechogenic endometrial transformation assessed on the day of hCG administration ($P < .001$).

Fanchin. Endometrial echogenicity and receptivity. Fertil Steril 1999.

RESEARCH

Open Access

Value of endometrial echo pattern transformation after hCG trigger in predicting IVF pregnancy outcome: a prospective cohort study

Zhaojuan Hou^{1,2}, Qiong Zhang^{1,2}, Jing Zhao^{1,2}, Aizhuang Xu^{1,2}, Aihua He^{1,2}, Xi Huang^{1,2}, Shi Xie^{1,2}, Jing Fu^{1,2}, Lan Xiao^{1,2} and Yanping Li^{1,2*}

- A total of 146 infertile women undergoing their first IVF cycle.
- A series of endometrial echo pattern monitoring was carried out in these patients after hCG trigger: hCG day, OPU + 1, OPU + 2, OPU + 3

Conclusions: The endometrial echogenicity value on OPU + 2 was recommended to evaluate endometrial receptivity. It seemed appropriate for clinicians to provide a ‘freeze all’ IVF cycle and transfer in a subsequent frozen-thawed embryos cycle when echogenicity value <60% on OPU + 2.

Table 2 Endometrial echogenicity values after hCG trigger in the two groups

Group	N	HCG	OPU + 1	OPU + 2	OPU + 3
Pregnant ^a	65	0.45 ± 0.14	0.75 ± 0.12	0.78 ± 0.12	0.81 ± 0.11
Non-pregnant	65	0.45 ± 0.17	0.67 ± 0.14	0.68 ± 0.14	0.72 ± 0.14
P		0.888	< 0.001	< 0.001	< 0.001

^aPatients who achieved a clinical pregnancy

ET LES DOSAGES HORMONAUX?

ESHRE guideline: ovarian stimulation for IVF/ICSI†

The ESHRE Guideline Group on Ovarian Stimulation, Ernesto Bosch¹, Simone Broer², Georg Griesinger³, Michael Grynberg⁴, Peter Humaidan⁵, Estratios Kolibianakis⁶, Michal Kunicki⁷, Antonio La Marca⁸, George Lainas⁹, Nathalie Le Clef¹⁰, Nathalie Massin¹¹, Sebastiaan Mastenbroek¹², Nikolaos Polyzos¹³, Sesh Kamal Sunkara¹⁴, Tanya Timeva¹⁵, Mira Töyli¹⁶, Janos Urbancsek¹⁷, Nathalie Vermeulen¹⁰, and Frank Broekmans^{2,*}

Recommandation basée sur Kwan et al. Cochrane 2014:

- 6 RCT → 781 femmes

Le dosage d'œstradiol ne semble pas améliorer les performances de l'écho pour prédire :

- Le risque OHSS
- Le nombre d'ovocytes prélevés
- Les chances de grossesse clinique

**Mais peut-être en pratique utile pour « ajuster » sa stimulation en cas d'échec...
mais pas vraiment d'études!!**

Recommendation

The addition of oestradiol measurements to ultrasound monitoring is probably not recommended.

Conditional ⊕⊕○○

Correlation between sonographic follow-up of follicular growth, serum and salivary estradiol in women undergoing controlled ovarian stimulation (IVF/ICSI)

A.S. ROTTIERS¹, L. DALEWYN¹, S. SOMERS¹, M.M. ALPER², D. SAKKAS², J. GERRIS¹

Facts Views Vis Obgyn 2018

- Strong correlation between serum and salivary E2 concentrations.
- In addition, both are strongly correlated with the product of the number of follicles and their average diameter, measured by TVUS.

421 RBMO VOLUME 42 ISSUE 2 2021

RBMO

ARTICLE

A multi-centre international study of salivary hormone oestradiol and progesterone measurements in ART monitoring

- Salivary estradiol hormone testing correlates well to serum-based assessment, whereas progesterone values, around the time of trigger, are not consistent from patient to patient.

Et la progestéronémie?

- Pas d'études évaluant « qualité de la stimulation ovarienne » et progestéronémie → pas recommandée pour évaluer ce paramètre!
- Mais impact d'une élévation prématurée de la progestérone en cours d'HSOC → ↓↓ taux d'implantation en cas de TEF
 - ✓ Incidence # 10 % des cycles (dépend du seuil choisi)
 - ✓ Publications récentes, seuil ≈ 1,5 ng/ml

Sonigo et al. RBMO 2014

Bosch et al, Hum Reprod 2010

Progesterone elevation and probability of pregnancy after IVF: a systematic review and meta-analysis of over 60 000 cycles

C.A. Venetis^{*}, E.M. Kolibianakis, J.K. Bosdou, and B.C. Tarlatzis

- Corrélation très nette entre élévation de progestérone J-hCG et baisse des taux de grossesse
- Problème des seuils différents au sein des études (techniques de dosages différentes +++)
- Importance du « terrain » = type de « répondeuse » en FIV

Table V Pooled ORs for pregnancy achievement in women with PE when compared with women without PE (per groups of PE threshold)

PE threshold group (ng/mL)	Achievement of pregnancy (ongoing pregnancy/live birth or clinical pregnancy)			Clinical pregnancy			Ongoing pregnancy/live birth		
	Number of studies or study subgroups/ total sample size	OR (95% CI)	Heterogeneity/ model	Number of studies or study subgroups/ total sample size	OR (95% CI)	Heterogeneity/ model	Number of studies or study subgroups/ total sample size	OR (95% CI)	Heterogeneity/ model
All studies									
0.4–0.6	5/1659	0.39 (0.14–1.08)	$P < 0.001$, $I^2 = 81.2\%/REM$	5/1659	0.37 (0.13–1.04)	$P < 0.001$, $I^2 = 81.2\%/REM$	2/1294	1.10 (0.67–1.81)	$P = 0.184$, $I^2 = 43.4\%/FEM$
0.8–1.1	40/16 304	0.79 (0.67–0.95)	$P < 0.001$, $I^2 = 66.8\%/REM$	40/16 304	0.79 (0.67–0.94)	$P < 0.001$, $I^2 = 67.3\%/REM$	15/7917	0.72 (0.56–0.94)	$P = 0.001$, $I^2 = 61.8\%/REM$
1.2–1.4	19/5885	0.67 (0.53–0.84)	$P = 0.003$, $I^2 = 53.4\%/REM$	14/2740	0.68 (0.49–0.94)	$P = 0.001$, $I^2 = 61.3\%/REM$	6/3316	0.64 (0.53–0.77)	$P = 0.49$, $I^2 = 0\%/FBM$
1.5–1.75	26/21 647	0.64 (0.54–0.76)	$P < 0.001$, $I^2 = 56.9\%/REM$	16/6785	0.65 (0.50–0.84)	$P < 0.001$, $I^2 = 64.5\%/REM$	14/ 15 971	0.62 (0.57–0.69)	$P = 0.54$, $I^2 = 0\%/FBM$
1.9–3.0	12/15 091	0.68 (0.51–0.91)	$P = 0.005$, $I^2 = 59.2\%/REM$	10/12 221	0.65 (0.41–1.04)	$P = 0.002$, $I^2 = 66.3\%/REM$	2/9447	0.67 (0.55–0.81)	$P = 0.69$, $I^2 = 0\%/FBM$

Analysis according to the type of population

EDITORIAL

Home monitoring of ovarian stimulation: an important step towards more patient-centred IVF

J.M.R. Gerris^{1,*}
B.C.J.M. Fauser²

TABLE 1 PUBLISHED STUDIES OF A STEP-WISE EXPLORATION OF THE CLINICAL APPLICABILITY OF PATIENT-OPERATED HOME SONOGRAPHY IN ART

Study reference	Key finding
<i>Gerris et al., 2009</i>	Patients are willing to use home sonography provided results of the treatment remain unaffected. (<i>n</i> = 25)
<i>Gerris et al., 2010</i>	Proof of concept in a hospital setting: patients are able to perform vaginal sonograms themselves. (<i>n</i> = 20)
<i>Gerris et al., 2014</i>	Randomized controlled trial: patient-operated home sonography gives similar results (embryological and clinical) to traditional sonographic follow up. Both patient-reported outcomes and cost are in favour of home sonography. (<i>n</i> = 121)
<i>Gerris et al., 2016</i>	Observational study of ICSI attempts using home sonography showed just one method failure. (<i>n</i> = 100)

COMMENTARY

Does artificial intelligence have a role in the IVF clinic?

Darren J X Chow^{1,2,3} , Philip Wijesinghe⁴ , Kishan Dholakia^{3,4,5,6} and Kylie R Dunning^{1,2,3}

Hurdles

- Most clinics are too small in isolation to have sufficient data for AI
- Limited computing power and data storage
- Discrepancy in recorded and reported patient characteristics and reproductive outcomes between clinics
- Variation between studies in the input parameters used for AI to predict reproductive outcomes

Requirements to develop AI in the infertility service

- Consortia to develop standardised and detailed recording and reporting of patient data
- Multi-clinic sharing of anonymised patient data
- Increased computing power and data storage
- Consensus on input parameters used in AI for predicting reproductive outcomes
- Rigorous, high quality, multi-centre, peer-reviewed validation studies for AI algorithms

REVIEW HYPOTHESIS

Artificial Intelligence in IVF: A Need

Charalampos Siristatidis^{1*}, Abraham Pouliakis², Charalampos Chrelas¹ and Dimitrios Kassanos¹

The neuron model

Multilayer neural networks

Table 1. Summary of features used for the prediction of IVF result.

Treatment Characteristics	Embryo Characteristics	Female Characteristics	Male Characteristics
Type of controlled ovarian hyperstimulation (i.e., long and short agonist, antagonist protocol)	Mean grade of embryos replaced	Female age	Male age
FSH dosage	Morphological scoring of human pronuclear zygotes	Infertility factor	Family history
Days of stimulation	Morphology score of the two best embryos available for transfer	Body Mass Index	Personal history
Basal E ₂ (pg/mL)	Cumulative embryo score	Mean ovarian volume (cm ³)	Sperm characteristics
Peak E ₂ level (pg/mL)	Number of embryos cleaved (number of embryos that have been divided)	Trauma	Sperm motility
Basal LH (mIU/mL)	Early cleavage morphology	Previous live births	Sperm function tests
Basal FSH (mIU/mL)	Early cleavage time	Previous abortions	Strict criteria for sperm morphology
Basal AFC	Equality of blastomeres	Duration of infertility	Chromatin packaging assessment
Inhibin B	Number of embryos	Diagnosis of infertility	Previous IVF cycles
Activin	Fragmentation rate	Smoking	Sperm DNA integrity assessment
AMH	Regularity of cells	Transfer Data	Hypoosmotic swelling test
CA-125	4-cell embryos at day 2	Type of catheter	Hemizona assay
Serum Vascular endothelial growth factor (VEGF)	Appearance of cytoplasm (darkness)	Transfer day	Zona pellucida -induced acrosome reaction
Challenge tests with clomiphene citrate	Freezing	Physician performing embryo transfer	
Ultrasonographic markers such as the AFC, total ovarian volume and blood flow	Prewash total motile sperm count	Embryo transfer technique	
3D ultrasonographic endometrial volume assessment on the day of hCG administration	Postwash total motile sperm count	Transfer with or without ultrasound guidance	
Endometrial thickness	Nucleus characteristics	Difficulty of transfer	
3D ultrasonography and power Doppler angiography	Visibility of nuclei in the cells	Type of fertilization (ICSI or IVF)	
Follicular fluid volume at egg recovery	Prewash total motile sperm count at egg recovery	Luteal support regimen	
Follicle size at egg recovery	Postwash total motile sperm count at egg recovery		
Number of eggs inseminated	Total number of embryos cultured		
Selection of euploid eggs	Number of embryos cultured on day 3		
Number of normal eggs fertilized	The ratio between the number of top embryos on day 3 and the total number of cultured embryos on day 3 (quality ratio)		

A machine learning algorithm can optimize the day of trigger to improve in vitro fertilization outcomes

Eduardo Hariton, M.D., M.B.A.,^a Ethan A. Chi,^b Gordon Chi,^b Jerrine R. Morris, M.D., M.P.H.,^a Jon Braatz, B.S.,^b Pranav Rajpurkar, Ph.D.,^b and Mitchell Rosen, M.D., H.C.L.D.^a

FIGURE 2

Feature importance bar chart. Removing a specific feature results in a drop in performance, with the most important features at the top, implying that those features are considered most when is the optimal day for trigger. It is important to note that feature importance is not designed to explain how important a given feature is to the outcome independently, but rather to explain our model's predictive capacity, as some features may be correlated. BMI = body mass index.

- Patients undergoing IVF-ICSI from 2008 to 2019 (n = 7,866).
 - Result(s): The machine learning algorithm is more effective than physician's decision :
 - ✓ 3.015 (95% CI 2.626, 3.371) more 2PNs.
 - ✓ 1.515 (95% CI 1.134, 1.871) more usable blastocysts.
- ➔ Future research is required to confirm these findings prospectively.

FIGURE 1

Points are colored red if a physician decided to trigger the given patient at that time and blue if otherwise.

Artificial intelligence in in vitro fertilization: a computer decision support system for day-to-day management of ovarian stimulation during in vitro fertilization

Gerard Letterie, M.D.,^a and Andrew Mac Donald, M.S.^b

An artificial intelligence platform to optimize workflow during ovarian stimulation and IVF: process improvement and outcome-based predictions

Gerard Letterie^{1,*}, Andrew MacDonald², Zhan Shi³

Conclusions : a first-iteration algorithm is described designed to improve workflow, minimize visits and level-load embryology work. This algorithm enables decisions at three interrelated nodal points for IVF workflow management to include monitoring on the single best day, assign trigger days to enable a range of 3 days for level-loading and estimate oocyte number.

FIGURE 1 Model for progressive data accumulation to predict outcomes and schedule the single best day, assign a trigger day and retrieval options and estimate the number of expected oocytes.

>10	-	0.84	0.76
Total retrieved mature oocytes	0.80	-	-
0-10	-	0.72	0.78
>10	-	0.78	0.81
Objective II: pre-IVF profile and best-day data			
Total retrieved oocytes	0.77	-	-
0-10	-	0.79	0.80
>10	-	0.74	0.74
Total retrieved mature oocytes	0.89	-	-
0-10	-	0.89	0.91
>10	-	0.88	0.86

Category I: pre-IVF profiles were defined as anti-Müllerian hormone concentration, age and body mass index prior to the start of stimulation.

Category II: pre-IVF profile and best-day data were defined as anti-Müllerian hormone concentration, age and body mass index plus oestradiol concentration and follicle number and diameters as measured on the predicted 'best day' or a single day of observation during ovarian stimulation.

AI in the treatment of fertility: key considerations

Jason Swain¹ · Matthew Tex VerMilyea² · Marcos Meseguer³ · Diego Ezcurra⁴ · Fertility AI Forum Group

Target	Objective	Company
Embryo assessment	EEVA viability status	https://hcp.merckgroup.com/en/fertility/technologies
	Geri Assess 2.0	https://hcp.merckgroup.com/en/fertility/technologies
	Viability and implantation potential	www.Presagen.com
	Viability and implantation potential	www.IVFBvision.ai
	Viability and implantation potential	www.FutureFertility.com
	Viability and implantation potential	www.Binflux.com
	Viability and implantation potential	www.IVY.ai
	Viability and implantation potential	STORK
	Viability and implantation potential	Fairtality
	Euploidy/aneuploidy status	www.Presagen.com
Oocyte assessment	Euploidy/aneuploidy status	ERICA— www.embryoranking.com
	Viability status	VIOLET— www.FutureFertility.com
Stimulation protocol	Predict live birth	www.UNIVFY.com
	Predict pregnancy	www.apricity.life
Uterine receptivity	Predict receptivity potential	www.FutureFertility.com

Conclusion

Intelligence Artificielle : des progrès majeurs en perspective !

Gain de temps

Efficiencce croissante

Optimiser la personnalisation de la prise en charge en AMP de chaque patiente / chaque couple

○

●